

lncRNA HAND2-AS1 Regulates Prostate Cancer Cell Growth Through Targeting the miR-106a-5p/RBM24 Axis

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Pengtao Wei¹
Jing Yang²
Dandan Zhang³
Meng Cui⁴
Lianjun Li^{5,6}

¹Department of Urology, Luoyang Central Hospital Affiliated to Zhengzhou University, Luoyang 471000, People's Republic of China; ²Central Sterile Supply Department, Shandong Provincial Hospital Affiliated to Shandong University, Jinan, People's Republic of China; ³Urinary Surgery, YiDu Central Hospital in Weifang City, Qingzhou, People's Republic of China; ⁴Department of Gynecology, Shandong Provincial Maternity and Childcare Hospital, Jinan, People's Republic of China; ⁵Department of Urology, Shandong Provincial Hospital Affiliated to Shandong University, Jinan 250014, People's Republic of China; ⁶Department of Urology, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan 250014, People's Republic of China

Introduction: Increasing evidence has shown that abnormally expressed long non-coding RNA (lncRNA) plays crucial roles in prostate cancer (PCa) progression.

Materials and Methods: Here, we analyzed the expression level of lncRNA HAND2 antisense RNA 1 (HAND2-AS1) in PCa cells and tissues. Function assays were performed to investigate the biological roles of HAND2-AS1 in PCa cell behaviors. Bioinformatics methods, luciferase activity reporter assay, and RNA pull-down assay were performed to validate the connection of microRNA-106a-5p (miR-106a-5p) with HAND2-AS1. Also, the target of miR-106a-5p was explored using the same methods.

Results: Our results revealed HAND2-AS1 expression was decreased in both PCa cells and tissues. In vitro functional assays showed HAND2-AS1 could inhibit PCa cell proliferation and colony formation through promoting cell apoptosis. Dual-luciferase activity assays showed miR-106a-5p could directly bind with HAND2-AS1 and RNA binding motif protein 24 (RBM24). Mechanistically, we showed that HAND2-AS1 regulates PCa cell behaviors via targeting miR-106a-5p/RBM24 axis.

Conclusion: In summary, our results illustrated that HAND2-AS1 functions as miR-106a-5p sponge to regulate RBM24 expression, and to influence PCa progression.

Keywords: HAND2-AS1, miR-106a-5p, RBM24, prostate cancer

Introduction

Surgery or androgen deprivation therapy can be used for prostate cancer (PCa) treatment for patients at early stages.¹ However, PCa remains a deadly malignancy for males in many countries and is even worse in developing countries.^{2,3} Therefore, it is essential to explore mechanisms underlying PCa progression to help better control PCa.

In recent years, many studies have indicated that long non-coding RNAs (lncRNAs) influence carcinogenesis.⁴ The improvements in high-throughput sequencing technology and bioinformatic analysis methods have identified numerous abnormally expressed lncRNAs in cancers.^{5,6}

HAND2 antisense RNA 1 (HAND2-AS1) is a lncRNA reported to function as tumor suppressive lncRNA in numerous cancers. For instance, HAND2-AS1 could serve as a sponge for microRNA-340-5p to regulate BCL2L1 expression in ovarian cancer.⁷ In esophageal squamous cell carcinoma, HAND2-AS1 overexpression could result in cancer cell proliferation, migration, and invasion inhibition by regulating miR-21.⁸ In chronic myeloid leukemia, HAND2-AS1 was shown to inhibit cancer

Correspondence: Lianjun Li
Department of Urology, Shandong Provincial Hospital Affiliated to Shandong University, Jinan 250014, P. R. China,
Department of Urology, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan 250014, People's Republic of China
Email lilianjun_jn@126.com

cell proliferation and promote apoptosis via regulating miR-1275.⁹ It is unclear whether HAND2-AS1 has a role in PCa.

miR-106a-5p is an miRNA reported to have decreased expression in renal cell carcinoma tissues and cells.¹⁰ It was found that the inhibition of miR-106a-5p expression could promote cancer cell migration and invasion through interacting with PAK5.¹⁰ On the other hand, elevated miR-106a-5p expression was identified in glioblastoma, and the miR-106a-5p overexpression could facilitate cancer invasion via regulating adenomatous polyposis coli.¹¹ These data appear to be contradictory as miR-106a-5p acted as oncogene in some cancers and tumor suppressor in others, which might be explained by the imperfect binding between miRNAs and targets.^{10,11} It has been revealed that RNA binding motif protein 24 (RBM24), which is an RNA binding protein, could directly interact with p21 and enhance the mRNA stability.¹² Moreover, RBM24 was revealed to have decreased expression in nasopharyngeal carcinoma and functioned as a direct target for miR-25 to inhibit cancer progression.¹³

In this study, we found decreased expression of HAND2-AS1 in PCa tissues and cells. Moreover, we showed HAND2-AS1 could regulate PCa cell proliferation, colony formation, and cell apoptosis in vitro. Moreover, the interaction of HAND2-AS1, miR-106a-5p, and RBM24 and their influences on PCa cell behaviors were further investigated to explore the mechanisms of HAND2-AS1 in PCa.

Materials and Methods

Cell Lines

PCa cells LNCaP, PC3 and DU145 and normal prostate epithelial cell P69 were acquired from Cell Bank of Chinese Academy of Science (Shanghai, China). Dulbecco's Modified Eagle's Medium (DMEM; Invitrogen, Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum (FBS; Invitrogen) was used to culture cells at a 37°C in a moist incubator containing 5% CO₂.

Cell Transfection

The full length of HAND2-AS1 or RBM24 was cloned into pcDNA3.1 by GenScript (Nanjing, Jiangsu, China). miR-106a-5p mimic, miR-106a-5p inhibitor and negative control (NC-mimic, or NC-inhibitor) were purchased from RiboBio (Guangzhou, Guangdong, China). Small interfering RNA against HAND2-AS1 (si-HAND2-AS1) and negative control (NC-siR) were also obtained from RiboBio. Cell transfection

was performed using Lipofectamine 2000 (Invitrogen). After 48 h of transfection, cells were collected for functional analyses.

Detection of HAND2-AS1, miR-106a-5p, and RBM24 Expression in Tumor Tissues and Normal Tissues

Expression levels of HAND2-AS1, miR-106a-5p, and RBM24 in PCa tissues and normal tissues were analyzed at ENCORI (<http://starbase.sysu.edu.cn/>).

Cell Proliferation Assay

Cells were plated in 96-well plates to analyze cell proliferation rate using cell counting kit-8 (CCK-8) method (Beyotime, Haimen, Jiangsu, China). CCK-8 reagent was added to the medium after seeding for 0, 1, 2, and 3 days. Then, optical density at the wavelength of 450 nm was measured using a microplate reader (Thermo Fisher Scientific, Inc.).

Colony Formation Assay

Cells were seeded in 6-well plates and incubated for 14 days to form colonies. Then, methanol was used to fix colonies, while crystal violet was used to stain these colonies. Finally, colony numbers were counted using a microscope.

Flow Cytometry Assay

Cells were collected, treated with trypsin, and stained with Annexin V-FITC and PI (Beyotime) in the dark following the supplier's instructions. Cell apoptosis rate was analyzed using flow cytometry (BD Biosciences, San Jose, CA, USA).

RNA Extraction and Quantitative Real-Time PCR (qRT-PCR)

RNA was extracted from cells using Trizol (Invitrogen) following the manufacturer's instructions. Using the PrimerScript kit, RNA was reverse transcribed into complementary DNA. qRT-PCR was performed with ABI 7500 (Applied Biosystems, Foster City, CA, USA) using SYBR Green (Takara, Dalian, Liaoning, China). Primers used in this work were as follows: HAND2-AS1: 5'-GGGTGTTTACG TAGACCAGAACC-3' (forward) and 5'-CTTCCAAAAG CCTTCTGCCTTAG-3' (reverse); RBM24: 5'-GGCCAAC GTGAACCTGGCATACTT-3' (forward) and 5'-GGCAGGT ATCCCGAAAGGTCTTTGT-3' (reverse); GAPDH: 5'-ATC ACTGCCACCCAGAAGAC-3' (forward) and 5'-TTTCTA GACGGCAGGTCAGG-3' (reverse); miR-106a-5p: 5'-GATGCTCAAAAAGTGCTTACAGTGCA-3' (forward)




Figure 1 LncRNA HAND2-AS1 was down-regulated in prostate cancer.

Notes: (A) The expression level of HAND2-AS1 was decreased in PCa tissues compared with adjacent tissues. (B) The expression level of HAND2-AS1 was down-regulated in PCa cell lines compared with normal cells. *** $P < 0.001$; ** $P < 0.01$.

Abbreviations: HAND2-AS1, HAND2 antisense RNA 1; PCa, prostate cancer.

and 5'-TATGGTTGTTCTGCTCTCTGTCTC-3' (reverse); U6 snRNA: 5'-AGAGCCTGTGGTGTCCG-3' (forward) and 5'-CATCTTCAAAGCACTTCCCT-3' (reverse). U6 snRNA was used as internal control for miR-106a-5p, while GAPDH was used as control for the rest genes.

Dual-Luciferase Reporter Assay

miR-106a-5p ranks top for all the targets for HAND2-AS1, while RBM24 ranks top among targets for miR-106a-4p. Sequence of HAND2-AS1 or RBM24 containing predicted miR-106a-5p binding sequence was inserted into Dual-luciferase Expression Vector (Promega, Madison, WI, USA) to generate HAND2-AS1 wild-type (HAND2-AS1-wt) or RBM24 wild-type (RBM24-wt). The mutant luciferase activity vector was built with/site-direct mutagenesis kit (Takara) and then these vectors were named HAND2-AS1-mt or RBM24-mt. After 48 h transfection, relative luciferase activities were measured using Dual-luciferase reporter assay system (Promega).

RNA Pull-Down Assay

The biotin labeled miR-106a-5p was synthesized by RiboBio. The biotin-miR-106a-5p was incubated with cell lysate, and then incubated with streptavidin magnetic agarose beads for 1 h. After washing the beads, RNA was isolated with Trizol and then subjected to qRT-PCR.

Statistical Analysis

Data were analyzed with GraphPad Prism 6.0 and then presented as mean \pm SD. Differences in groups were analyzed using Student's *t*-test or one-way analysis of

variance (ANOVA) and Tukey post hoc test. *P* value less than 0.05 indicated significant difference.

Results

HAND2-AS1 Expression Was Decreased in PCa Tissues and Cells

Expression level of HAND2-AS1 in PCa tissues and normal tissues was analyzed with ENCORI and we found HAND2-AS1 expression was decreased in PCa tissues compared with normal tissues (Figure 1A). Meanwhile, qRT-PCR results indicated that HAND2-AS1 expression was lower in PCa cells than in normal cell line (Figure 1B).

HAND2-AS1 Overexpression Inhibited PCa Cell Growth

HAND2-AS1 expression level was elevated by pHAND2-AS1 transfection (Figure 2A). CCK-8 assay indicated HAND2-AS1 overexpression inhibited cell proliferation (Figure 2B) and the colony formation assay suggested that HAND2-AS1 overexpression inhibited colony formation (Figure 2C). Moreover, flow cytometry analysis revealed that pHAND2-AS1 transfection promoted cell apoptosis (Figure 2D).

HAND2-AS1 Knockdown Promoted PCa Cell Growth

qRT-PCR showed that introduction of si-HAND2-AS1 decreased the levels of HAND2-AS1 in PCa cells (Figure 3A). CCK-8 assay, colony formation assay,




Figure 2 LncRNA HAND2-AS1 overexpression inhibited PCa.

Notes: (A) pHAND2-AS1 transfection increased HAND2-AS1 level in PCa cells. (B) CCK-8 assay showed pHAND2-AS1 group inhibited PCa cells' proliferation. (C) Colony formation number in cells with pHAND2-AS1 transfection. (D) Cell apoptosis rate of pHAND2-AS1 group increased. *** $P < 0.001$; ** $P < 0.01$.

Abbreviations: HAND2-AS1, HAND2 antisense RNA 1; PCa, prostate cancer; CCK-8, cell counting kit-8.

and flow cytometry assay indicated that knockdown of HAND2-AS1 promoted cell growth (Figure 3B-D).

HAND2-AS1 Functioned as a Sponge for miR-106a-5p

The binding module of HAND2-AS1 and miR-106a-5p was presented in Figure 4A. Luciferase activity reporter assay indicated that miR-106a-5p overexpression inhibited luciferase activity in cells with HAND2-AS1-wt transfection (Figure 4B). RNA pull-down assay indicated the direct interaction of HADN2-AS1 and miR-106a-5p (Figure 4C). Moreover, we showed that miR-106a-5p expression level

was increased in PCa tissues and cells (Figure 4D and 4E). qRT-PCR showed that pHAND2-AS1 transfection decreased miR-106a-5p expression, while miR-106a-5p mimic transfection increased miR-106a-5p expression (Figure 4F). Functional assays showed groups with miR-106a-5p mimic transfection exhibited higher cell growth ability compared with the other groups (Figure 4G-I).

HAND2-AS1 Served as a Sponge of miR-106a-5p and Regulated RBM24

Bioinformatic tool showed RBM24 was a possible target for miR-106-5p, as shown in Figure 5A. Luciferase




Figure 3 LncRNA HAND2-AS1 knockdown promoted PCa.

Notes: (A) si-HAND2-AS1 transfection decreased HAND2-AS1 level in PCa cells. (B) CCK-8 assay showed si-HAND2-AS1 group increased PCa cells' proliferation. (C) Colony formation number in cells with si-HAND2-AS1 transfection. (D) Cell apoptosis rate of si-HAND2-AS1 group decreased. *** P < 0.001.

Abbreviations: HAND2-AS1, HAND2 antisense RNA 1; PCa, prostate cancer; CCK-8, cell counting kit-8; si-HAND2-AS1, small interfering RNA against HAND2-AS1; NC-siR, negative control small interfering RNA.

intensity in PCa cells was significantly reduced after co-transfection of RBM24-wt and miR-106a-5p mimic compared with that of RBM24-wt and NC-mimic (Figure 5B). Exploration of RBM24 expression level in PCa showed RBM24 levels were significantly decreased in PCa tissues and cells (Figure 5C and D). Results of qRT-PCR showed RBM24 expression level was stimulated by pRBM24 but inhibited by miR-106a-5p mimic (Figure 5E). Functional assays showed cell growth ability was highest in groups with miR-106a-5p mimic transfection but lowest in those with pRBM24 transfection (Figure 5F-H).

Discussion

Numerous lncRNAs have been identified as aberrantly expressed in PCa and participating in PCa progression.¹⁴ For example, Jiang et al revealed lncRNA NEAT1 was expressed at a high level in docetaxel-resistant PCa cells and exerted an oncogenic effect by regulating ACSL4 expression through sponging both miR-34a-5p and miR-204-5p.¹⁵ Moreover, lncRNA CASC15 was also found to be expressed at a high level in PCa tissues and cells.¹⁶ Moreover, the knockdown of CASC15 was found to suppress PCa cell migration and invasion through functioning as a sponge for miR-200a-3p.¹⁶





Figure 4 HAND2-AS1 served as a sponge of miR-106a-5p.

Notes: (A) The targeting relationship between HAND2-AS1 and miR-106a-5p. (B) The luciferase intensity of PCa cells transfected with HAND2-AS1-wt/HAND2-AS1-mt and miR-106a-5p mimic and NC-mimic. (C) RNA pull-down assay to demonstrate the correlation of HAND2-AS1 and miR-106a-5p. (D) The expression level of miR-106a-5p was increased in PCa tissues compared with adjacent tissues. (E) The expression level of miR-106a-5p was upregulated in PCa cell lines compared with normal cells. (F) miR-106a-5p expression in PCa cells transfected with pcDNA3.1+miR-106a-5p mimic, pHAND2-AS1+miR-106a-5p mimic, pHAND2-AS1+NC-mimic, pcDNA3.1+NC-mimic, pcDNA3.1+miR-106a-5p inhibitor, and pcDNA3.1+NC-inhibitor. (G) CCK-8 assay showed cell proliferation rate in cells transfected with pcDNA3.1+miR-106a-5p mimic, pHAND2-AS1+miR-106a-5p mimic, pHAND2-AS1+NC-mimic, pcDNA3.1+NC-mimic, pcDNA3.1+miR-106a-5p inhibitor, and pcDNA3.1+NC-inhibitor. (H) Colony formation assay showed colony numbers in cells transfected with pcDNA3.1+miR-106a-5p mimic, pHAND2-AS1+miR-106a-5p mimic, pHAND2-AS1+NC-mimic, pcDNA3.1+NC-mimic, pcDNA3.1+miR-106a-5p inhibitor, and pcDNA3.1+NC-inhibitor. (I) Flow cytometry showed cell apoptosis rate in cells transfected with pcDNA3.1+miR-106a-5p mimic, pHAND2-AS1+miR-106a-5p mimic, pHAND2-AS1+NC-mimic, pcDNA3.1+NC-mimic, pcDNA3.1+miR-106a-5p inhibitor, and pcDNA3.1+NC-inhibitor. *** P < 0.001; ** P < 0.01; * P < 0.05.

Abbreviations: HAND2-AS1, HAND2 antisense RNA 1; PCa, prostate cancer; CCK-8, cell counting kit-8; miR-106a-5p, microRNA-106a-5p; NC-mimic, negative control for miR-106a-5p mimic; wt, wild-type; mt, mutant.

OncoTargets and Therapy downloaded from https://www.dovepress.com/ by 202.120.23.46 on 19-Jun-2020
 For personal use only.




Figure 5 miR-106a-5p regulated expression of RBM24.

Notes: (A) The targeting relationship between RBM24 and miR-106a-5p. (B) The luciferase intensity of PCa cells transfected with RBM24-wt/RBM24-mt and miR-106a-5p mimic and NC-mimic. (C) The expression level of RBM24 was decreased in PCa tissues compared with adjacent tissues. (D) The expression level of RBM24 was downregulated in PCa cell lines compared with normal cells. (E) RBM24 expression in PCa cells transfected with pcDNA3.1+miR-106a-5p mimic, pRBM24+miR-106a-5p mimic, pRBM24+NC-mimic, pcDNA3.1+NC-mimic, pHAND2-AS1+miR-106a-5p mimic, and pHAND2-AS1+NC-mimic. (F) CCK-8 assay showed cell proliferation rate in cells transfected with pcDNA3.1+miR-106a-5p mimic, pRBM24+miR-106a-5p mimic, pRBM24+NC-mimic, and pcDNA3.1+NC-mimic. (G) Colony formation assay showed colony numbers in cells transfected with pcDNA3.1+miR-106a-5p mimic, pRBM24+miR-106a-5p mimic, pRBM24+NC-mimic, and pcDNA3.1+NC-mimic. (H) Flow cytometry showed cell apoptosis rate in cells transfected with pcDNA3.1+miR-106a-5p mimic, pRBM24+miR-106a-5p mimic, pRBM24+NC-mimic, and pcDNA3.1+NC-mimic. *** P < 0.001; ** P < 0.01; * P < 0.05.

Abbreviations: RBM24, RNA binding motif protein 24; PCa, prostate cancer; CCK-8, cell counting kit-8; miR-106a-5p, microRNA-106a-5p; NC-mimic, negative control for miR-106a-5p mimic; wt, wild-type; mt, mutant.

In this work, we first validated the downregulated status of HAND2-AS1 in both PCa tissues and cells. Gain-of and loss-of experiments indicated that HAND2-AS1 overexpression inhibits PCa cell proliferation, colony formation, and promotes apoptosis, whereas knockdown of HAND2-AS1 exerted opposite effects on PCa cell

behaviors. These results suggested a tumor suppressive role of HAND2-AS1 in PCa, which is consistent with its role in cancers including ovarian cancer, esophageal squamous cell carcinoma, and chronic myeloid leukemia.⁷⁻⁹ In recent years, the competing endogenous RNA (ceRNA) theory was used to understand the functions of lncRNA.¹⁷

The connection of HAND2-AS1 and miR-106a-5p was validated by luciferase activity reporter assay and RNA pull-down assay. Here, we showed miR-106a-5p overexpression could stimulate – while miR-106a-5p knockdown could inhibit – PCa cell growth. These results indicated that miR-106a-5p serves as an oncomiR in PCa, which is consistent with its role in glioblastoma.¹¹ Moreover, rescue experiments showed that miR-106a-5p was a functional target of HAND2-AS1. In addition, we found RBM24 was a putative target for miR-106a-5p, and the overexpression of RBM24 decreases PCa cell growth, suggesting its tumor suppressive role in PCa. In this work, the functional analyses revealed that lncRNA HAND2-AS1 exerted a tumor suppressive effect in PCa by regulating RBM24 expression through serving as a sponge for miR-106a-5p.

Conclusion

To sum up, we identified that HAND2-AS1 was remarkably decreased in PCa tissues and cells. Besides, HAND2-AS1 could inhibit PCa cell proliferation and colony formation by promoting apoptosis through downregulating RBM24 via sponging miR-106a-5p. These results suggested that HAND2-AS1 functioned as a tumor suppressive lncRNA in PCa and may be a potential therapeutic target for PCa.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Zhu Y, Yang XQ, Han CT, et al. Pathological features of localized prostate cancer in China: a contemporary analysis of radical prostatectomy specimens. *PLoS One*. 2015;10(3):e121076.
- Litwin MS, Tan HJ. The diagnosis and treatment of prostate cancer: a review. *JAMA*. 2017;317(24):2532–2542. doi:10.1001/jama.2017.7248
- Guo T, Wang XX, Fu H, Tang YC, Meng BQ, Chen CH. Early diagnostic role of PSA combined miR155 detection in prostate cancer. *Eur Rev Med Pharmacol Sci*. 2018;22(6):1615–1621. doi:10.26355/eurrev_201803_14568
- Ponting CP, Oliver PL, Reik W. Evolution and functions of long noncoding RNAs. *Cell*. 2009;136(4):629–641. doi:10.1016/j.cell.2009.02.006
- Zhao X, Tang DY, Zuo X, Zhang TD, Wang C. Identification of lncRNA-miRNA-mRNA regulatory network associated with epithelial ovarian cancer cisplatin-resistant. *J Cell Physiol*. 2019;234(11):19886–19894. doi:10.1002/jcp.28587
- Wang B, Tang D, Zhang Z, Wang Z. Identification of aberrantly expressed lncRNA and the associated TF-mRNA network in hepatocellular carcinoma. *J Cell Biochem*. 2020;121(2):1491–1503. doi:10.1002/jcb.29384
- Chen J, Lin Y, Jia Y, Xu T, Wu F, Jin Y. LncRNA HAND2-AS1 exerts anti-oncogenic effects on ovarian cancer via restoration of BCL2L11 as a sponge of microRNA-340-5p. *J Cell Physiol*. 2019;234(12):23421–23436. doi:10.1002/jcp.28911
- Yan Y, Li S, Wang S, et al. Long noncoding RNA HAND2-AS1 inhibits cancer cell proliferation, migration, and invasion in esophagus squamous cell carcinoma by regulating microRNA-21. *J Cell Biochem*. 2019;120(6):9564–9571. doi:10.1002/jcb.28233
- Yang JR, Shi MX, Zeng Y. LncRNA HAND2-AS1 inhibits proliferation and promotes apoptosis of chronic myeloid leukemia cells by sponging with miRNA-1275. *Eur Rev Med Pharmacol Sci*. 2019;23(5):2103–2111. doi:10.26355/eurrev_201903_17254
- Pan Y-J, Wei -L-L, Wu X-J, Huo F-C, Mou J, Pei D-S. MiR-106a-5p inhibits the cell migration and invasion of renal cell carcinoma through targeting PAK5. *Cell Death Dis*. 2017;8(10):e3155. doi:10.1038/cddis.2017.561
- Li D, Wang Z, Chen Z, et al. MicroRNA-106a-5p facilitates human glioblastoma cell proliferation and invasion by targeting adenomatous polyposis coli protein. *Biochem Biophys Res Commun*. 2016;481(3–4):245–250. doi:10.1016/j.bbrc.2016.10.132
- Jiang Y, Zhang M, Qian Y, Xu E, Zhang J, Chen X. Rbm24, an RNA-binding protein and a target of p53, regulates p21 expression via mRNA stability. *J Biol Chem*. 2014;289(6):3164–3175. doi:10.1074/jbc.M113.524413
- Hua WF, Zhong Q, Xia TL, et al. RBM24 suppresses cancer progression by upregulating miR-25 to target MALAT1 in nasopharyngeal carcinoma. *Cell Death Dis*. 2016;7(9):e2352. doi:10.1038/cddis.2016.252
- Hua JT, Chen S, He HH. Landscape of noncoding RNA in prostate cancer. *Trends Genet*. 2019;35(11):840–851. doi:10.1016/j.tig.2019.08.004
- Jiang X, Guo S, Zhang Y, et al. LncRNA NEAT1 promotes docetaxel resistance in prostate cancer by regulating ACSL4 via sponging miR-34a-5p and miR-204-5p. *Cell Signal*. 2019;65:109422. doi:10.1016/j.celsig.2019.109422
- Zhang C, Wang GX, Fu B, Zhou XC, Li Y, Li YY. LncRNA CASC15 promotes migration and invasion in prostate cancer via targeting miR-200a-3p. *Eur Rev Med Pharmacol Sci*. 2019;23(19):8303–8309. doi:10.26355/eurrev_201910_19141
- Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the Rosetta stone of a hidden RNA language? *Cell*. 2011;146(3):353–358. doi:10.1016/j.cell.2011.07.014

OncoTargets and Therapy

Dovepress

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>